



Variational iteration method for twelfth-order boundary-value problems

A.S.V. Ravi Kanth*, K. Aruna

Applied Mathematics Division, School of Science and Humanities, V.I.T. University, Vellore – 632 014, Tamil Nadu, India

ARTICLE INFO

Keywords:

Variational iteration method
Linear–nonlinear problems
Boundary-value problems
Integral equations

ABSTRACT

In this paper, He's variational iteration method is applied to solve twelfth-order boundary-value problems. The numerical results obtained with minimum amount of computation are compared with the exact solutions to show the efficiency of the method. The results show that the variational iteration method is of high accuracy, more convenient and efficient for solving high-order boundary-value problems.

© 2009 Elsevier Ltd. All rights reserved.

1. Introduction

We consider the twelfth-order boundary-value problem of the type:

$$y^{xii}(x) + f(x)y = g(x) \quad a < x < b \quad (1)$$

$$y^{2n}(0) = \alpha_{2n}, \quad y^{2n}(1) = \beta_{2n}, \quad n = 0, 1, \dots, 5 \quad (2)$$

where α_{2n} and β_{2n} ($n = 0, 1, \dots, 5$) are finite real constants and the functions $f(x)$ and $g(x)$ are continuous on $[a, b]$. This type of boundary-value problem arises in an infinite horizontal layer of fluid is heated from below, with the assumption that a uniform magnetic field is also applied across the fluid in the same direction as gravity and the fluid is subject to the action of rotation, instability sets in. When instability sets in as ordinary convection, it is modeled by tenth-order boundary-value problems, when instability sets in as over stability, it is modeled by twelfth-order boundary-value problems (see Chandrasekhar [1]). Agarwal's book [2] contains theorems which detail the conditions for existence and uniqueness of solutions of the twelfth-order boundary-value problems. Different numerical methods have been proposed by various authors for twelfth-order boundary-value problems. To mention a few, finite difference method [3,4], modified decomposition method [5], twelfth degree splines [6] and thirteen degree splines [7].

In this paper, we aim to apply the variational iteration method proposed by He [8–14] to twelfth-order boundary-value problems. This method has been employed to solve a large variety of linear and nonlinear problems such as the KdV, the K(2, 2), the Burgers and the cubic Boussinesq equations [23], Korteweg–de Vries (KdV) and the modified Korteweg–de Vries equations [16], nonlinear differential equations of fractional order [17], KdV Burgers equations and Lax's seventh-order KdV equations [18], generalized Burgers–Fisher and Burgers equations [19], Helmholtz equation [20], semi-linear inverse parabolic equation [15], linear fractional differential equation [21], nonlinear partial differential equations [22] etc.

2. Variational iteration method

According to the variational iteration method [8–14], we can construct a correction functional for twelfth-order boundary-value problem (1)–(2) in the form

$$y_{n+1}(x) = y_n(x) + \int_0^x \lambda [y_n^{xii}(s) + f(s)y_n(s) - g(s)] ds \quad (3)$$

* Corresponding author.

E-mail address: asvrvikanth@yahoo.com (A.S.V. Ravi Kanth).

where λ is a general Lagrange multiplier which can be identify via variational theory [8–10]. Making $y_{n+1}(x)$ stationary with respect to $y_n(x)$, we can identify the Lagrange multiplier, which reads

$$\lambda = \frac{(s-x)^{11}}{11!}. \quad (4)$$

Example 1. First, we consider the following BVP [7]

$$y^{xii}(x) + xy(x) + (120 + 23x + x^3)e^x = 0 \quad 0 < x < 1 \quad (5)$$

with boundary conditions

$$\left. \begin{aligned} y(0) = 0 = y(1), y'(0) = 1, y'(1) = -e, y''(0) = 0, \\ y''(1) = -4e, y'''(0) = -3, y'''(1) = -9e, y^{iv}(0) = -8, \\ y^{iv}(1) = -16e, y^v(0) = -15, y^v(1) = -25e \end{aligned} \right\}. \quad (6)$$

According to (3), we have the following iteration formulation

$$y_{n+1}(x) = y_n(x) + \int_0^x \frac{(s-x)^{11}}{11!} [y_n^{xii}(s) + sy_n(s) + (120 + 23s + s^3)e^s] ds. \quad (7)$$

Now, we assume that an initial approximation has the form

$$y_0(x) = a + bx + cx^2 + dx^3 + ex^4 + fx^5 + gx^6 + hx^7 + ix^8 + jx^9 + kx^{10} + lx^{11} \quad (8)$$

where $a, b, c, d, e, f, g, h, i, j, k$ and l are unknown constants to be further determined. By iteration formula (7), we have the following first-order approximation

$$\begin{aligned} y_1(x) &= y_0(x) + \int_0^x \frac{(s-x)^{11}}{11!} [y_0^{xii}(s) + sy_0(s) + (120 + 23s + s^3)e^s] ds \\ &= -\frac{l}{1295295050649600}x^{24} - \frac{k}{647647525324800}x^{23} - \frac{j}{309744468633600}x^{22} \\ &\quad - \frac{i}{140792940288000}x^{21} - \frac{h}{60339831552000}x^{20} - \frac{g}{24135932620800}x^{19} \\ &\quad - \frac{f}{8892185702400}x^{18} - \frac{e}{2964061900800}x^{17} - \frac{d}{871782912000}x^{16} - \frac{c}{217945728000}x^{15} \\ &\quad - \frac{b}{43589145600}x^{14} - \frac{a}{6227020800}x^{13} + \left(l + \frac{13}{5702400}\right)x^{11} + \left(k + \frac{1}{72576}\right)x^{10} \\ &\quad + \left(j - \frac{1}{40320}\right)x^9 + \left(i - \frac{23}{10080}\right)x^8 + \left(h - \frac{41}{1008}\right)x^7 + \left(g - \frac{59}{120}\right)x^6 + \left(f - \frac{109}{24}\right)x^5 \\ &\quad + \left(e - \frac{98}{3}\right)x^4 + \left(d - \frac{359}{2}\right)x^3 + (c - 715)x^2 + (b - 1849)x + (a - 2340) \\ &\quad + (2340 - 491x + 36x^2 - x^3)e^x. \end{aligned}$$

Incorporating the boundary conditions, Eq. (6), into $y_1(x)$, and solving the system simultaneously, we obtain

$$\begin{aligned} a &= 0, & b &= 1, & c &= 0, & d &= -0.5, & e &= -1/3, & f &= -1/8, & g &= -0.03333333362911, \\ h &= -0.00694444319430, & i &= -0.00119047835708, & j &= -0.00017360919905, \\ k &= -0.00002204671105, & l &= -0.00000248000384. \end{aligned}$$

We, therefore obtain the following first-order approximate solution

$$\begin{aligned} y_1(x) &= \frac{0.0000024899384}{1295295050649600}x^{24} + \frac{0.00002204671105}{647647525324800}x^{23} \\ &\quad + \frac{0.00017360919905}{309744468633600}x^{22} + \frac{0.001190478357105}{140792940288000}x^{21} + \frac{0.00694444319430}{60339831552000}x^{20} \\ &\quad + \frac{0.03333333362911}{24135932620800}x^{19} + \frac{1/8}{8892185702400}x^{18} + \frac{1/3}{2964061900800}x^{17} \\ &\quad + \frac{1/2}{87178291200}x^{16} - \frac{1}{43589145600}x^{14} - 2.00261976924803e-007x^{11} \end{aligned}$$

Table 1

Comparison of the first-order approximate solution, Eq. (9), with the exact solution.

x	VIM solution $y_1(x)$	Exact solution	Absolute error
0.0	0	0	0
0.1	0.09946538262776	0.09946538262681	9.524186994624984e–013
0.2	0.19542444130575	0.19542444130563	1.245392677873269e–013
0.3	0.28347034959130	0.28347034959096	3.351208199831035e–013
0.4	0.35803792743337	0.35803792743390	5.379585665821196e–013
0.5	0.41218031767423	0.41218031767503	8.044676036433884e–013
0.6	0.43730851209258	0.43730851209372	1.140698646651117e–012
0.7	0.42288806856919	0.42288806856880	3.933520176246930e–013
0.8	0.35608654855867	0.35608654855879	1.23068222796986e–013
0.9	0.22136428000330	0.22136428000413	8.254508188088039e–013
1.0	3.268502189e–013	0	3.268502189413308e–013

$$\begin{aligned}
& -8.268051438007055e - 006x^{10} - 1.984107863515873e - 004x^9 \\
& - 0.00347222438883x^8 - 0.04761904636890x^7 - 0.52500000029578x^6 \\
& - 4.66666666666667x^5 - 33x^4 - 180x^3 - 715x^2 - 1848x - 2340 \\
& + (2340 - 491x + 36x^2 - x^3)e^x.
\end{aligned} \tag{9}$$

The exact solution of the above differential equations is

$$y(x) = x(1 - x)e^x.$$

Comparison of the first-order approximate solution with the exact solution is tabulated in Table 1 showing a remarkable agreement.

Example 2. Next, we consider the following BVP [7]

$$y^{xii}(x) - y(x) + 12(2x \cos x + 11 \sin x) = 0 \quad -1 < x < 1 \tag{10}$$

$$y(-1) = 0 = y(1), \quad y'(-1) = y'(1) = 2 \sin(1),$$

$$y''(-1) = -y''(1) = -4 \cos(1) - 2 \sin(1),$$

$$y^{(iii)}(-1) = y^{(iii)}(1) = 6 \cos(1) - 6 \sin(1),$$

$$y^{(vi)}(-1) = -y^{(iv)}(1) = 8 \cos(1) + 12 \sin(1),$$

$$y^{(v)}(-1) = y^{(v)}(1) = -20 \cos(1) + 10 \sin(1). \tag{11}$$

According to (3), we have the following iteration formulation

$$y_{n+1}(x) = y_n(x) + \int_0^x \frac{(s-x)^{11}}{11!} [y_n^{xii}(s) - y_n(s) + 12(2s \cos s + 11 \sin s)] ds. \tag{12}$$

Now, we assume that an initial approximation has the form

$$y_0(x) = a + bx + cx^2 + dx^3 + ex^4 + fx^5 + gx^6 + hx^7 + ix^8 + jx^9 + kx^{10} + lx^{11} \tag{13}$$

where $a, b, c, d, e, f, g, h, i, j, k$ and l are unknown constants to be further determined. By iteration formula (12), we have the following first-order approximation

$$\begin{aligned}
y_1(x) = & \frac{l}{647647525324800}x^{23} + \frac{k}{309744468633600}x^{22} + \frac{j}{140792940288000}x^{21} \\
& + \frac{i}{60339831552000}x^{20} + \frac{h}{24135932620800}x^{19} + \frac{g}{8892185702400}x^{18} \\
& + \frac{f}{2964061900800}x^{17} + \frac{e}{871782912000}x^{16} + \frac{d}{217945728000}x^{15} \\
& + \frac{c}{43589145600}x^{14} + \frac{b}{6227020800}x^{13} + \frac{a}{479001600}x^{12} \\
& + \left(l - \frac{1}{369600}\right)x^{11} + \left(j + \frac{1}{6048}\right)x^9 + \left(h - \frac{1}{420}\right)x^7 + \left(f - \frac{3}{10}\right)x^5 + a \\
& + (b - 132)x + cx^2 + (d + 14)x^3 + ex^4 + gx^6 + ix^8 + kx^{10} + 156 \sin x - 24x \cos x.
\end{aligned}$$

Incorporating the boundary conditions, Eq. (11), into $y_1(x)$, and solving the system simultaneously, we obtain

$$a = -0.277555755576842759e - 16, \quad b = -0.9999999999999996891,$$

Table 2a

Comparison of the first-order approximate solution, Eq. (14), with the exact solution.

x	Exact solution	VIM solution $y_1(x)$	Absolute error
−1.0	0	−0.932587340685e−14	0.93258734068513e−14
−0.9	0.1488321128292218	0.1488321128292236	0.1776356839400e−14
−0.8	0.2582481927238280	0.2582481927238152	0.1282307593442e−13
−0.7	0.3285510204912224	0.3285510204912206	0.1831867990631e−14
−0.6	0.3613711829728227	0.3613711829728373	0.1465494392505e−13
−0.5	0.3595691539531523	0.3595691539531660	0.1376676550535e−13
−0.4	0.3271114075392664	0.3271114075392686	0.1332267629550e−14
−0.3	0.2689233880618189	0.2689233880618338	0.1482147737874e−13
−0.2	0.1907225575632587	0.1907225575632596	0.9436895709313e−15
−0.1	0.0988350824803598	0.0988350824803585	0.5551115123125e−16
0	0	−.277555755576e−16	0.27755575557684e−16

Table 2b

Comparison of the first-order approximate solution, Eq. (14), with the exact solution.

x	Exact solution	VIM solution $y_1(x)$	Absolute error
0.0	0	−2.77555755576e−017	2.7755575557684e−017
0.1	−0.0988350824803598	−0.0988350824803584	1.3877787807814e−015
0.2	−0.1907225575632587	−0.1907225575632595	8.3266726846886e−016
0.3	−0.2689233880618190	−0.2689233880618338	1.6542323066914e−014
0.4	−0.3271114075392664	−0.3271114075392686	2.1649348980190e−015
0.5	−0.3595691539531523	−0.3595691539531660	1.3766765505351e−014
0.6	−0.3613711829728227	−0.3613711829728373	2.2204460492503e−015
0.7	−0.3285510204912224	−0.3285510204912206	1.6098233857064e−014
0.8	−0.2582481927238281	−0.2582481927238155	1.2601031329495e−014
0.9	−0.1488321128292218	−0.1488321128292238	1.9984014443252e−015
1.0	0	0.9325873406851e−14	9.3258734068513e−015

$$\begin{aligned}
c &= -0.971445158559381916e-16, & d &= 1.1666666666666667, \\
e &= 0.100845239619727242e-15, & f &= -0.175000000000001599, \\
g &= -0.345504587738423279e-16, & h &= 0.853174603174756963e-2, \\
i &= 0.955915575530589730e-17, & j &= -0.201168430335752597e-3, \\
k &= -0.132768981572782405e-17, & l &= 0.278078403089638659e-5.
\end{aligned}$$

We, therefore obtain the following first-order approximate solution

$$\begin{aligned}
y_1(x) = & \frac{0.278078403089638659e-5}{647647525324800}x^{23} - \frac{0.132768981572782405e-17}{309744468633600}x^{22} \\
& - \frac{0.201168430335752597e-3}{140792940288000}x^{21} + \frac{0.955915575530589730e-17}{60339831552000}x^{20} \\
& + \frac{0.853174603174756963e-2}{24135932620800}x^{19} - \frac{0.345504587738423279e-16}{8892185702400}x^{18} \\
& - \frac{0.175000000000001599}{2964061900800}x^{17} + \frac{0.100845239619727242e-15}{871782912000}x^{16} \\
& + \frac{1.1666666666666667}{217945728000}x^{15} - \frac{0.971445158559381916e-16}{43589145600}x^{14} \\
& - \frac{0.99999999999996891}{6227020800}x^{13} - \frac{0.277555755576842759e-16}{479001600}x^{12} \\
& + 0.751563252686809709e-7x^{11} - 0.451134989999223829e-4x^9 \\
& + 0.615079365079518846e-2x^7 - 0.475000000000001587x^5 \\
& - 0.277555755576842759e-16 - 133x - 0.971445158559381916e-16x^2 \\
& + 15.1666666666666667x^3 + 0.100845239619727242e-15x^4 \\
& - 0.3455045877384279e-16x^6 + 0.955915575530589730e-17x^8 \\
& - 0.132768981572782405e-17x^{10} + 156 \sin x - 24x \cos x.
\end{aligned} \tag{14}$$

The exact solution of the above differential equations is $y(x) = (x^2 - 1) \sin x$. Comparison of the first-order approximate solution with the exact solution is tabulated in Tables 2a and 2b showing a remarkable agreement.

3. Conclusion

Variational iteration method is remarkably effective for solving twelfth-order boundary-value problems. One iteration is enough to obtain very high accurate solution. This method is very powerful, efficient and promoting, which will be certainly found wide applications.

References

- [1] S. Chandrasekhar, Hydrodynamic and Hydromagnetic stability, Dover, New York, 1981.
- [2] R.P. Agrawal, Boundary Value Problems for Higher Ordinary Differential Equations, World Scientific, Singapore, 1986.
- [3] A. Boutayeb, E.H. Twizell, Finite difference methods for twelfth order boundary value problems, J. Comput. Appl. Math. 35 (1991) 133–138.
- [4] E.H. Twizell, A. Boutayeb, K. Djidjedi, Numerical methods for eight-, tenth-, twelfth- order eigenvalue problems arising in thermal instability, Adv. Comput. Math. 2 (1994) 407–436.
- [5] A.M. Wazwaz, Approximate solutions to boundary value problems of higher order by the modified decomposition method, Comput. Math. Appl. 40 (2000) 679–691.
- [6] S.S. Siddiqi, E.H. Twizell, Spline solutions of linear twelfth order boundary value problems, J. Comput. Appl. Math. 78 (1997) 371–390.
- [7] S.S. Siddiqi, A. Ghazala, Solutions of twelfth order boundary value problems using thirteen degree spline, Appl. Math. Comput. 182 (2006) 1443–1453.
- [8] J.H. He, Variational iteration method: A kind of non linear analytical technique: Some examples, Internat. J. Nonlinear Mech. 244 (1999) 699–708.
- [9] J.H. He, Variational iteration method for autonomous ordinary differential systems, Appl. Math. Comput. 114 (2000) 115–123.
- [10] J.H. He, Some asymptotic methods for strongly non linear equations, Internat. J. Modern Phys. B. 20 (2006) 1141–1199.
- [11] J.H. He, Variational principle for some nonlinear partial differential equations with variable coefficients, Chaos Soliton Fractals 19 (2004) 847–851.
- [12] J.H. He, The variational iteration method for eighth-order initial-boundary value problems, Phys. Scr. 76 (2007) 680–682.
- [13] J.H. He, X.H. Wu, Variational iteration method: New development and applications, Comput. Math. Appl. 54 (2007) 881–894.
- [14] J.H. He, Variational iteration method — Some recent results and new interpretations, J. Comput. Appl. Math. 207 (2007) 3–17.
- [15] Mehdi Tatari, Mehdi Dehghan, He's variational iteration method for computing a control parameter in a semi-linear inverse parabolic equation, Chaos Solitons Fractals 33 (2007) 671–677.
- [16] M. Inc, Numerical simulation of KdV and mKdV equations with initial conditions by the variational iteration method, Chaos Solitons Fractals 34 (2007) 1075–1081.
- [17] Z.M. Odibat, S. Momani, Application of variational iteration method to nonlinear differential equations of fractional order, Int. J. Nonlinear Sci. Numer. Simul. 7 (2006) 27–34.
- [18] A.A. Soliman, A numerical simulation and explicit solutions of KdV–Burgers' and Lax's seventh-order KdV equations, Chaos Solitons Fractals 29 (2006) 294–302.
- [19] Mahdi Moghimi, Fatemeh S.A Hejazi, Variational iteration method for solving generalized Burger–Fisher and Burger equations, Chaos Solitons Fractals 33 (2007) 1756–1761.
- [20] S. Momani, S. Abuasad, Application of He's variational iteration method to Helmholtz equation, Chaos Solitons Fractals 27 (2006) 1119–1123.
- [21] Shaher Momani, Zaid Odibat, Numerical comparison of methods for solving linear differential equations of fractional order, Chaos Solitons Fractals 31 (2007) 1248–1255.
- [22] N. Bildik, A. Konuralp, The use of variational iteration method, differential transform method and adomian decomposition method for solving different types of nonlinear partial differential equations, Int. J. Nonlinear Sci. Numer. Simul. 7 (2006) 65–70.
- [23] A.M. Wazwaz, The variational iteration method for rational solutions for KdV, K(2, 2), Burgers, and cubic Boussinesq equations, J. Comput. Appl. Math. 207 (2007) 18–23.